

# AVALIAÇÃO CLÍNICA DA EFICIÊNCIA DE BARREIRA ANTISSEPTICA MEDICAMENTOSA NO CONTROLE DE CONTAMINAÇÃO DO SÍTIO REGENERADO DURANTE AS EXPOSIÇÕES DE MEMBRANA, EM PROCEDIMENTOS DE RGT.

**OBJETIVO:** Verificação da eficiência de barreira antisseptica no controle da contaminação do sítio a ser regenerado, nas exposições da membrana.

**METODOLOGIA:** Mínimo 30 casos ou sítios.

## ***Grupo teste***

- 1 - Após a exposição induzida nas regiões de dentes extraídos ou outras situações, fazer a barreira antisséptica com Alvogyl colocando-o sob os bordos da ferida. Ver livro RGT.
- 2 - Fazer a limpeza com spray e a troca da medicação a cada 7 dias.
- 3 - Utilizar os colutórios regularmente – clorexidina 0,12%.
- 4- Verificar a cor, morfologia dos tecidos externos e internos, i.e., sob a membrana. Após a remoção da membrana, verificar a contaminação bacteriana ou não, no lado interno. Pode-se optar por cultura microbiológica ou exame direto visual com microscopia de luz.

## ***Grupo Controle***

- 1 - Não aplicar a barreira antisséptica.
- 2 - Usar o colutório.
- 3 - Verificar as condições ao remover a membrana.

## **BIBLIOGRAFIA:**

Al Shawi A. Open-packing method for the everely comminuted fractured mandible due to missile injury. Br J Oral Maxillofac Surg 1995; 33(1): 36-39.

Anderson M, Seow WK. pH of endodontic medicaments used in pediatric dentistry: effects of dyadic combinations. *J Clin Pediatr Dent* 1990; 15(1): 42-45.

Bramante CM, Berbert A. Influence of time of calcium hydroxide iodiform paste replacement in the treatment root perforations. *Braz Dent J* 1994;1: 45-51.

Callskan MK, Piskin B. International resorption after accidental extrusion of iodoform past into the mandibular canal. *Endod Dent Traumatol* 1993; 9(2): 81-84.

Chindia ML. A conservative management of an extensive odontogenic residual cyst: a case report. *East Afr Med J* 1991; 68(2): 143-148.

Cruz M. *Regeneração Guiada Tecidual*. São Paulo: Santos Editora; 2006.

Garth RJ, Brightwell AP. A comparison of packing materials used in nasal surgery. *J Laryngol Otol* 1994;8(7):564-566.

Harry P, Bouachor G, Tirot P, Chennebaut JM, Cailleux A, Allain P, Alquier P. Iodoform poisoning. 3 cases. *J Toxicol Clin Exp* 1992; 12(1):35-42.

Holan G, Fuks AB. A comparison of pulpectomies using ZOE and KRI past in primary molars: a retrospective study. *Pediatr Dent* 1993; 15(6): 403-407.

Kawakami T, Nakamura C, Hasegawa H, Akahane S, Eda S. Ultrastructural study of initial calcification in the rat subcutaneous tissue elicited by root canal filling material. *Oral Surg Oral Med Pathol* 1987; 63(3):360-365.

Kawakami T, Nakamura C, Eda S. Effects of the penetration of a root filling material into the mandibular canal .2. Changes in the alveolar nerve tissue. *Endod Dent Traumatol* 1991; 7(1):42-47.

Lyll JB. Third molar surgery: The effect of primary closure, wound dressing and metronidazole on postoperative recovery. *JR Army Med Corps* 1991; 137(2): 100-103.

MSDS for Iodoform, Infosafe no: AJ1CF, 17/02/92, Alax Chemicals.

Persson LG, Lekholm U, Leonhardt A, Dahlen G, Lindhe J. Bacterial colonization on internal surfaces of Branemark system implant components. *Clin Oral Implants Res* 1996; 7(2):90-95.

Saggar V, Chandra S, Jaiswal JN, Singh M. Antimicrobial efficacy of iodoformized zinc oxide-eugenol sealer on microorganisms of root canal. *J Indian Soc Pedod Prev Dent* 1991;1:1-3.

Shipochliev T, Dimitrov A, Aleksandrova E. Anti-inflammatory action for a group of plant extracts. *Vet Med Nauki* 1981; 18(6): 87-94.

Syrjanen SM, Syrjanen KJ. Influence of Alvogyl on the healing of extraction wound in man. *Int J Oral Surg* 1979; 8(1): 22-30.

Syrjanen SM, Syrjanen KJ. Plasma cells and their Immunoglobulins in the normal and delayed healing of the extraction wound in man. *Br J Oral Surg* 1980;18(2): 100-106.

Szakiel A, Wasiukiewicz I, Janiszowska W. Metabolism for [3-3H] oleanolic in the Isoletd *Calendula officialis* leaf cells and transport of glycosides to the cell wall and the. *Acta Biochim Pol* 1995; 42(1): 25-29.

Thomas AM, Chandra S, Chandra S, Pandey RK. Elimination of infection in pulpectomized deciduous teeth: a short-term study using iodoform paste. *J Endod* 1994; 20(5): 233-235.

Wright KJ, Barbosa SV, Araki K, Spangberg LS. In vitro antimicrobial and cytotoxic effects of Kri 1 paste and zinc oxide-eugenol used in primary tooth pulpectomies. *Pediatr Dent* 1994; 16(2):102-106.